

A service of the National Library of Medicine
and the National Institutes of Health

My NCBI [?] [Sign In] [Register]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for Go Clear

Limits Preview/Index History Clipboard Details

Display Abstract Show 20 Sort by Send to

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorials

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI

Related Resources

Order Documents

NLM Mobile

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

1: J Androl. 1989 Jul-Aug;10(4):270-4.

Related Articles, Links

The effect of tumor necrosis factor on human sperm motility in vitro.**Eisermann J, Register KB, Strickler RC, Collins JL.**

Department of Obstetrics and Gynecology, Washington University School of Medicine, St. Louis, Missouri 63110.

Tumor necrosis factor (TNF alpha) is present in elevated levels in peritoneal fluid from infertile women with endometriosis. The effect of TNF alpha on human sperm motility in vitro was evaluated utilizing peritoneal fluid from infertile women with minimal endometriosis containing 0, 100, 400, or 800 U of TNF alpha/ml as well as similar concentrations of recombinant human TNF alpha. No reduction in progressive and total motility was found at recombinant TNF alpha concentrations of 100 U/ml. However, 500 and 1000 U of recombinant TNF alpha/ml caused a significant reduction in progressive and total sperm motility after 4 and 21 hours of incubation when compared with controls. Similarly, peritoneal fluid containing 100 U of TNF alpha/ml did not significantly reduce progressive and total sperm motility after either 4 or 21 hours of incubation; but peritoneal fluid containing 400 U of TNF alpha/ml reduced progressive sperm motility after 4 and 21 hours and total sperm motility after 21 hours of incubation. Peritoneal fluid with a TNF alpha concentration of 800 U/ml caused a significant reduction in both progressive and total sperm motility after 4 and 21 hours when compared with controls of TNF alpha-negative peritoneal fluid. The addition of polyclonal rabbit anti-TNF alpha antibody or 30-min heat inactivation at 56 C of TNF alpha-positive peritoneal fluid reversed the inhibitory effect on sperm motility. The ability of TNF alpha to cause a significant reduction of sperm motility in vitro suggests that this may be a mechanism for the infertility observed in women with minimal endometriosis.

PMID: 2777718 [PubMed - indexed for MEDLINE]

Display Abstract Show 20 Sort by Send to

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

Jun 6 2006 06:32:22

KARGER

Login / Subscriptions: **US Patent & Trademark Office**

Gynecologic and Obstetric Investigation

Journal H-

■ Journal C-

Guideline:

Editorial E

Aims and

Subscripti

Download

Vol. 53, Suppl. 1, 2002

[Free Abstract](#) [Article \(References\)](#) [Article \(PDF 207 KB\)](#)

The Fourth Japan Conference on Endometriosis
Current Perspectives on Endometriosis and Infertility
Proceedings of the Fourth Japan Conference on Endometriosis
Hakone Prince Hotel, Kanagawa, Japan, April 28, 2001
Guest Editors: H. Hoshiai, Osaka-Sayama; Y. Taketani, Tokyo; N. Terakawa,
Yonago

Session 2: Chairman: Hoshiai, H. (Osaka-Sayama). Original Article

Role of Cytokines in Endometriosis-Associated Infertility

Tomio Iwabe, Tasuku Harada, Naoki Terakawa

Department of Obstetrics and Gynecology, Tottori University School of Medicine,
Yonago, Japan

[Address of Corresponding Author](#)

Gynecologic and Obstetric Investigation 2002;53:19-25 (DOI: 10.1159/000049420)



Key Words

- Endometriosis
- Peritoneal fluid
- Interleukin-6
- Infertility



Abstract

Endometriosis, which is common in women of reproductive age, may affect fertility. It is also clear that mechanical disruption of the pelvic anatomy may cause infertility. However, our understanding of the association between the early stage of endometriosis and infertility remains incomplete. Bloody peritoneal fluid (PF) is

frequently observed in the cul-de-sac of endometriosis patients and contains various biologically active factors. We found that the concentrations of tumor necrosis factor alpha (TNF- α) and interleukin-6 (IL-6) in PF from patients with endometriosis were significantly higher than that of patients with endometriosis. There were significantly positive correlations between the levels of TNF- α and IL-6. We compared the levels of these cytokines with regard to the R-AFS stages and scores, but no differences were observed. In contrast, these cytokines correlate with the number and extent of red color peritoneal endometriosis. TNF- α increased the expression of IL-6 messenger RNA and protein in endometriotic stromal cells derived from chocolate cyst in a dose-dependent manner. The addition of IL-6 inhibited the development of mouse preimplantation embryo and impaired sperm function. We concluded that increased levels of IL-6 in peritoneal fluid of patients with active red endometriosis might be related to endometriosis-associated fertility.

Copyright © 2002 S. Karger AG, Basel



Author Contacts


Tomio Iwabe, MD
Department of Obstetrics and Gynecology
Tottori University School of Medicine
36-1, Nishimachi, Yonago, 683-8504 (Japan)
Tel. +81 859 34 8127, Fax +81 859 34 8089, E-Mail iwabe@biken.osaka-u.ac.jp



Article Information

Number of Print Pages : 7
Number of Figures : 6, Number of Tables : 2, Number of References : 7

[Free Abstract](#) [Article \(References\)](#) [Article \(PDF 207 KB\)](#)

 Copyright © 2006 S. Karger AG, Basel



US 20020064826A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2002/0064826 A1**
Ruben et al. (43) **Pub. Date: May 30, 2002**

(54) **CYTOKINE RECEPTOR-LIKE
POLYNUCLEOTIDES, POLYPEPTIDES, AND
ANTIBODIES**

provisional of provisional application No. 60/168,
621, filed on Dec. 3, 1999.

Publication Classification

(76) Inventors: **Steven M. Ruben**, Olney, MD (US);
Jian Ni, Germantown, MD (US); **Paul**
E. Young, Gaithersburg, MD (US);
Yanggu Shi, Gaithersburg, MD (US)

(51) **Int. Cl.⁷** **C07H 21/02**; C07H 21/04;
C12P 21/02; C12N 5/06
(52) **U.S. Cl.** **435/69.1**; 435/325; 435/320.1;
536/23.1

Correspondence Address:

HUMAN GENOME SCIENCES INC
9410 KEY WEST AVENUE
ROCKVILLE, MD 20850

(57) **ABSTRACT**

(21) Appl. No.: **09/874,069**

(22) Filed: **Jun. 6, 2001**

Related U.S. Application Data

(63) Continuation-in-part of application No. PCT/US00/
32525, filed on Nov. 30, 2000, which is a non-

The present invention relates to novel human cytokine receptor-like polypeptides and isolated nucleic acids containing the coding regions of the genes encoding such polypeptides. Also provided are vectors, host cells, antibodies, and recombinant methods for producing human cytokine receptor-like polypeptides. The invention further relates to diagnostic and therapeutic methods useful for diagnosing and treating disorders related to these novel human cytokine receptor-like polypeptides.



A service of the National Library of Medicine
and the National Institutes of Health

www.pubmed.gov

My NCBI
[Sign In] [Register]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed

for sperm cytokine

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorials

New/Noteworthy

E-Utilities

Note: Performing your original search, *sperm cytokine*, in PubMed will retrieve **485** citations.

Display Abstract Show 20 Sort by Send to

All: 1 Review: 0

1: Arch Androl. 1995 Nov-Dec;35(3):173-85.

[Related Articles](#), [Links](#)

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI

Effect of cytokines on sperm motility and ionophore-stimulated acrosome reaction.

Fedder J, Ellerman-Eriksen S.

Department of Gynecology and Obstetrics, University Hospital of Aarhus, Denmark.

The effects of soluble products from leukocytes (WBC) on sperm motility and calcium ionophore-stimulated acrosome reaction (AR) were examined. Supernatants of mononuclear WBC, isolated from peripheral blood and stimulated by the lectins *Phytolacca americana* (pokeweed mitogen) or concanavalin A, caused a weak but significant inhibition of progressive sperm motility. The recombinant cytokine interferon-gamma (IFN-gamma) in high concentrations inhibited motility of sperm from 5 of 8 donors tested. The recombinant cytokines tumor necrosis factor-alpha (TNF-alpha) and interleukin-8 (IL-8) did not show any effect on sperm motility. Reactive oxygen species (ROS) produced by stimulation of polymorphonuclear WBC with a phorbol ester (PMA) tended to inhibit sperm motility. Neither supernatants from lectin-stimulated mononuclear WBC, IFN-gamma, TNF-alpha, IL-8, nor ROS showed any significant effect on the ionophore-challenged AR.

PMID: 8585773 [PubMed - indexed for MEDLINE]

Display Abstract Show 20 Sort by Send to

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Jun 6 2006 06:32:22